

Revised: August 2017 (8th version)

Standard Commodity Classification No. of Japan
875200

- Kampo-preparation -

# TEIKOKU Saikokaryukotsuboreito Extract Granules

&lt; Saikokaryukotsuboreito &gt;

<b>Storage</b>
Store at room temperature in a tight container. [See the "Precaution for handling" section.]

Approval No.	(61AM) 3675
Date of listing in the NHI reimbursement price	October 1987
Date of initial marketing in Japan	October 1987

<b>Expiration date</b>
Saikokaryukotsuboreito should be used before the expiration date indicated on the label and the package.

## DESCRIPTION

Composition	9.0 g of TEIKOKU Saikokaryukotsuboreito Extract Granules contains 4.33 g of a dried water extract of the following mixed crude drugs.	
	JP Bupleurum Root -----	5.0 g
	JP Pinellia Tuber -----	4.0 g
	JP Poria Sclerotium -----	3.0 g
	JP Cinnamon Bark -----	3.0 g
	JP Scutellaria Root -----	2.5 g
	JP Jujube -----	2.5 g
	JP Ginger -----	1.0 g
	JP Ginseng -----	2.5 g
	JP Longgu -----	2.5 g
	JP Oyster Shell -----	2.5 g
	JP Rhubarb -----	1.0 g
	(JP: The Japanese Pharmacopoeia)	
	Inactive ingredients	JP Lactose Hydrate JP Microcrystalline Cellulose JP Magnesium Stearate
Description	Dosage form	Granules
	Color	Light brown
	Smell	Slightly characteristic smell
	Taste	Bitter
	ID code	TEIKOKU 12

## PRECAUTIONS

### 1. Careful Administration (Saikokaryukotsuboreito should be administered with care in the following patients.)

- (1) Patients with diarrhea or soft feces [These symptoms may be aggravated.]
- (2) Patients with an extremely weak gastrointestinal tract [Anorexia, epigastric distress, nausea, abdominal pain, diarrhea, etc. may occur.]
- (3) Patients with greatly declined constitution [Adverse reactions are likely to occur, and the symptoms may be aggravated.]

### 2. Important Precautions

- (1) When this product is used, the patient's "SHO" (constitution/symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- (2) When this product is coadministered with other Kampo-preparations (Japanese traditional herbal medicines), etc., attention should be paid to the duplication of the contained crude drugs. Special caution should be exercised when this product is coadministered with preparations containing Rhubarb.
- (3) Since there is an individual difference in the cathartic action of Rhubarb, caution should be exercised concerning the dosage and administration.

SHO : The term "SHO" refers to a particular pathological status of a patient evaluated by the Kampo diagnosis, and is patterned according to the patient's constitution, symptoms, etc. Kampo-preparations (Japanese traditional herbal medicines) should be used after confirmation that it is suitable for the identified "SHO" of the patient.

## INDICATIONS

The following symptoms of those patients with palpitation and insomnia who have mental instability:

Symptoms associated with hypertension (palpitation, anxiety, and insomnia), neurosis, climacteric neurosis, and night cry in children

## DOSAGE AND ADMINISTRATION

For oral use, the usual adult dosage is 3.0 g of TEIKOKU Saikokaryukotsuboreito Extract Granules three times daily before meals.

The dosage may be adjusted according to the patient's age, body weight, and symptoms.

### 3. Adverse Reactions

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

#### (1) Clinically significant adverse reactions

**1) Interstitial pneumonia:** If fever, cough, dyspnea, abnormal pulmonary sound (fine crackle), etc. are observed, administration of this product should be discontinued, and examinations such as X-ray should be performed immediately and appropriate measures such as administration of adrenocortical hormones taken. Besides, the patient should be advised to discontinue this product immediately and to make contact with the physician in the event of fever, cough, dyspnea, etc.

**2) Hepatic dysfunction and jaundice:** Hepatic dysfunction and/or jaundice with elevation of AST (GOT), ALT (GPT), Al-P and  $\gamma$ -GTP or other symptoms may occur. The patient should be carefully monitored for abnormal findings. Administration should be discontinued and appropriate therapeutic measures should be taken, if abnormalities are observed.

#### (2) Other adverse reactions

	Incidence unknown
<b>Hypersensitivity</b> <sup>Note)</sup>	Rash, Redness, Pururitus, Urticaria, etc.
<b>Gastrointestinal</b>	Anorexia, Epigastric distress, Abdominal pain, Diarrhea, etc.

Note) If such symptoms are observed, administration should be discontinued.

### 4. Use in the Elderly

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

### 5. Use during Pregnancy, Delivery or Lactation

(1) Use of this product in pregnant women, women who may possibly be pregnant is not recommended. [The uterotonic action and congestive action on the intrapelvic organs of Rhubarb contained in this product may cause premature birth or abortion.]

(2) This product should be administered with care in nursing mothers. [Anthraquinone derivatives in Rhubarb contained in this product may be excreted in breast milk and induce diarrhea in nursing infants.]

### 6. Pediatric Use

The safety of this product in children has not been established. [Insufficient clinical data.]

### PRECAUTIONS FOR HANDLING

1. This product should not be stored in direct rays and should be stored in a cool place with, if possible, little humidity.
2. Since it is hygroscopic property, this product should not be stored in humid places after opening.

### PACKAGING

Bottles of 500 g

Boxes of 3.0 g× 42 packets

Boxes of 3.0 g×252 packets

### REQUEST FOR LITERATURE SHOULD BE MADE TO:

Product Information Unit,  
Pharmaceutical Sales & Marketing Dept.  
Teikoku Seiyaku Co., Ltd.  
6-6 Nihonbashi-kobunacho, Chuo-Ku, Tokyo  
103-0024, Japan  
Tel 0120-189-567

### Manufactured by:

Teikoku Kampo Seiyaku Co., Ltd.  
80-11 Kitahara, Donari, Donari-cho, Awa, Tokushima  
771-1506, Japan

### Distributed by:

Teikoku Seiyaku Co., Ltd.  
567, Sanbonmatsu, Higashikagawa, Kagawa  
769-2695, Japan